

FOR EPILEPSY*

SPRITAM[®]

Newest dosage form of levetiracetam

Easy-to-take SPRITAM provides the safety profile and seizure reduction of levetiracetam[†]

**A SMALL CHANGE CAN
MAKE A BIG DIFFERENCE**

Not an actual patient.

***Indications and Usage**

SPRITAM (levetiracetam) is a prescription medicine taken by mouth that is used to treat partial-onset seizures in people 4 years of age and older. SPRITAM is also used with other medicines to treat myoclonic seizures in people 12 years of age and older with juvenile myoclonic epilepsy, and primary generalized tonic-clonic seizures in people 6 years of age and older with certain types of generalized epilepsy. SPRITAM is recommended for use in people weighing more than 20 kg (44 lbs).

[†]Based on comparable rate and extent of absorption to levetiracetam immediate release tablets.

Please see additional Important Safety Information inside, and enclosed Full Prescribing Information and Medication Guide.

SAY GOODBYE TO MESSY MEASURING AND BULKY PILLS

WHEN AN EASY-TO-TAKE FORM MATTERS

A typical young person with childhood epilepsy will take thousands of doses of their seizure medication in their lifetime.

THOUSANDS
OF DOSES



Important Safety Information

SPRITAM may not be for everyone. Ask your healthcare provider if SPRITAM is right for you. Do not take SPRITAM if you are allergic to levetiracetam.



SAY HELLO TO SPRITAM[®]

Spritam[®]
(levetiracetam)
Tablets for Oral Suspension

SPRITAM IS THE UNIQUE, FLASH-DISPERSING TECHNOLOGY

At home or on the go, **SPRITAM**, the newest dosage form of levetiracetam, simplifies administration with its unique flash-dispersing technology. Surprisingly easy to give and easy-to-take, **SPRITAM** is gone in a flash with a sip of liquid. Seize the opportunity to simplify the daily epilepsy medication routine with **SPRITAM**.

Warnings and Precautions

Antiepileptic drugs, including SPRITAM, may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. Call your healthcare provider right away if you have new or worsening symptoms of depression, any unusual changes in mood or behavior, or suicidal thoughts, behavior, or thoughts about self-harm that you have never had before or may be worse than before.

SPRITAM may cause extreme sleepiness, tiredness, and weakness, and problems with muscle coordination. You should not drive, operate machinery or do other dangerous activities until you know how SPRITAM affects you.

*SPRITAM disintegrates in a mean time of 11 seconds (range of 2 to 27 seconds).
Important Safety Information continued on next page



SPRITAM'S unique flash-dispersing technology disintegrates quickly in your mouth with a sip of liquid*, delivering an exact dose every time.



TAKE AN ACTIVE ROLE IN MANAGING EPILEPSY

WHAT KIND OF MEDICATION ROUTINE WOULD YOU PREFER?

- I want to feel in control and confident that the exact dose is administered every time
- I want to look forward to family activities free of discomfort at medication administration time
- I want a medication that fits our lifestyle with a dosing plan we can sustain

Warnings and Precautions – continued

SPRITAM can cause severe allergic reactions after the first dose or at any time during treatment. Discontinue SPRITAM and seek immediate medical attention if you have allergic reactions such as swelling of the face, lips, eyes, tongue and throat, trouble swallowing or breathing, and hives.



TALK TO YOUR DOCTOR ABOUT SPRITAM

A FEW QUESTIONS TO ANSWER BEFORE YOUR NEXT APPOINTMENT

- Are you having problems giving or taking your current medication?
- What caused the last missed dose?
- Do you have difficulty measuring your medication?
- Now that you've experienced a placebo demonstrator tablet, how might this dosage form fit in with your lifestyle?

Warnings and Precautions – continued

Serious skin rashes can happen after you start taking SPRITAM. There is no way to tell if a mild rash will become a serious reaction. Call your healthcare provider right away at the first sign of rash.

Do not stop taking SPRITAM unless instructed by your healthcare provider. Stopping a seizure medication suddenly can cause you to have seizures more often or seizures that will not stop, which is a very serious problem.

If you become pregnant or intend to become pregnant, contact your healthcare provider right away.

SPRITAM SUPPORT



YOU MAY BE ELIGIBLE FOR SAVINGS ON SPRITAM

Visit [SPRITAM.COM/SAVINGS](https://www.spritam.com/savings) for details.
When prescribed, be sure to ask for
SPRITAM by name at your pharmacy.

Common Adverse Reactions

In clinical trials, the most common side effects seen in adults who take SPRITAM include sleepiness, weakness, dizziness, and infection. In addition to those previously listed, the most common side effects seen in children who take SPRITAM include tiredness, acting aggressive, nasal congestion, decreased appetite, and irritability.



Common Adverse Reactions – continued

Talk to your healthcare provider about other possible side effects with SPRITAM. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**. To report SUSPECTED ADVERSE REACTIONS, contact Aprecia Pharmaceuticals, LLC at **1-844-882-7732** or FDA at **1-800-FDA-1088** or www.fda.gov/medwatch.

Administration

SPRITAM is intended to disintegrate in the mouth when taken with a sip of liquid. As a main way to take SPRITAM, place a tablet on the tongue with a dry hand, and then take a sip of liquid. SPRITAM should be swallowed only after the tablet disintegrates in the mouth.

SPRITAM should not be swallowed intact. Partial tablet(s) should not be administered.

Alternately, add a whole SPRITAM tablet(s) to a small volume of liquid in a cup and swirl gently. Swallow right after the tablet disintegrates. If there is any medicine left in the cup, rinse with a small volume of liquid, swirl and swallow the remaining contents.

For additional safety information, please see US Full Prescribing Information and Medication Guide. This information does not take the place of talking with your healthcare provider about your condition or your treatment.



A small change can make a big difference



**TALK TO YOUR DOCTOR
ABOUT SPRITAM**



Manufactured by Aprecia Pharmaceuticals, LLC, Blue Ash, OH 45242 USA. Distributed by Prasco, LLC, Mason, OH 45040 USA. SPRITAM® is a registered trademark of Aprecia Pharmaceuticals, LLC.

© 2022 Prasco LLC all rights reserved. PP-MA-SP-US-0297